

## PATHOTYPIC DIVERSITY OF *Rhynchosporium secalis* ISOLATES IN FIVE PROVINCES OF IRAN \*

S. BEIGI<sup>1\*\*</sup>, H. ZAMANIZADEH<sup>1</sup>, M. RAZAVI<sup>2</sup> and R. ZARE<sup>2</sup>

(Received: 11. 12. 2010; Accepted: 20. 7. 2011)

### Abstract

Leaf blotch of barley, caused by *Rhynchosporium secalis*, is a major disease of barley in Iran. Pathogenic variation of barley scald pathogen was determined by testing 47 isolates collected from five provinces of Iran. Among the isolates tested Rs55 and Rs86 were the most virulent and Rs45 was the least virulent. According to their reactions against eight differential barley cultivars containing known resistance genes, 20 different pathotypes were determined and these pathotypes represent a very broad virulence spectrum. Pathotype 1 (pt1) was found in all five surveyed provinces. Pathotypes pt15, pt12 and pt13 caused susceptible reaction in 62, 50 and 50% of cultivars, respectively. Of these cultivars Igri and Armele containing *BRR4* and *BRR1* resistance genes were the most resistant whereas cultivars WI and Digger were the most susceptible against these pathotypes, respectively. There was no association between pathogenic variability and geographical location of *R. secalis*.

**Keywords:** Barley, Scald, Differential cultivars, Resistance genes, Aggressiveness.

See Persian text for figures and tables (Pages ۳۳۱ –۳۴۰).

---

\*: A Part of PhD. Thesis of the First Author, Submitted to College of Agriculture and Natural Resource, Islamic Azad University, Science and Research Branch, Tehran, Iran.

\*\* : Corresponding Author, Email: siamakb\_30@yahoo.com

1. PhD. Student and Prof. of Plant Pathology, College of Agriculture and Natural Resource, Islamic Azad University, Branch of Science and Research, Tehran, Iran.

2. Assis. Prof. and Prof. Iranian Research Institute of Plant Protection, Tehran, Iran.

## References

- ALI, S.M., MAYFIELD, A.H. and CLARE, B.G. 1976. Pathogenicity of 203 isolates of *Rhynchosporium secalis* on 21 barley cultivars. **Plant Pathol.** 9:135-143.
- ARABI, M.L.E., JAWAHR, M. and AL-SHEHADEH, E. 2008. Molecular and pathogenic variation identified among isolates of *Rhynchosporium secalis*. **J. Plant Pathol.** 90: 179-184.
- BOUAJILA, A., ABANG, M.A., HAOUAS, S., UDUPA, S., REZGUI, S., BAUM, M. and YAHYAOU, A. 2007. Genetic diversity of *Rhynchosporium secalis* in Tunisia as revealed by pathotype, AFLP, and microsatellite analyses. **Mycopathologia** 163: 281-294.
- COOKE, L.R., LOCKE, T. and LOCKLEY, K.D. 2004. The effect of fungicide programmes based on proconazole on the control and DMI sensitivity of *Rhynchosporium secalis* in winter barley. **Crop Protec.** 23: 393-406.
- GOODWIN, S.B., ALLARD, R.W. and WEBSTER, R.K. 1990. A nomenclature for *Rhynchosporium secalis* pathotypes. **Phytopathology** 80: 1330-1336.
- GOODWIN, S.B., ALLARD, R.W., HARDY, S.A. and WEBSTER, R.K. 1992. Hierarchical structure of pathogenic variation among *Rhynchosporium secalis* populations in Idaho and Oregon. **Can. J. Bot.** 70: 810-17.
- GOODWIN, S.B., WEBSTER, R.K. and ALLARD, R.W. 1994. Evidence for mutation and migration as sources of genetic variation in population of *Rhynchosporium secalis*. **Phytopathology** 94: 1047-1053.
- JACKSON, L.F., KAHLER, A.L., WEBSTER, R.K. and ALLARD, R.W. 1978. Conservation of scald resistance in barley composite cross population. **Phytopathology** 68: 645-650.
- JORGENSEN, H.J.L. and SMEDEGARD-PETERSEN, V. 1995. Pathogenic variation of *Rhynchosporium secalis* in Denmark and sources of resistance in barley. **Plant Dis.** 79: 297-301.
- KARI, A.G. and GRIFFITHS, E. 1993. Components of partial resistance of barley to *Rhynchosporium secalis*: use of seedling tests to predict field resistance. **Ann. Appl. Biol.** 123: 545-561.
- MATHER, D.E. 1985. Compendium of Barley Diseases. APS Press. 249pp.
- McDERMOTT, J.M., McDONALD, B.A. ALLARD, R.W. and WEBSTER, R.K. 1989. Genetic variability for pathogenicity, isozyme, ribosomal DNA and colony color variants in populations of *Rhynchosporium secalis*. **Genetics** 122: 551-565.
- McDONALD, B.A. 1989. The population biology of host-pathogen interactions. **Ann. Rev. Phytopathol.** 27: 77-94.
- MELES, K., HULLUKA, M. and ABANG. M.M. 2004. Phenotypic diversity in *Rhynchosporium secalis* from Ethiopia and host response to barley scald. **Plant Pathol. J.** 3: 26-34.
- NEWTON, A.C., SEARLE, J., GUY, D.C., HACKETT, C.A. and COOKE, D.E.L. 2001. Variability in pathotype, aggressiveness, RAPD profile, and rDNA ITS1 sequences of UK isolates of *Rhynchosporium secalis*. **J. Plant Dis.** 108: 446-458.
- OXLEY, S.J.P., COOKE, L.R., BLACK, L.A. and MERCER, P.C. 2003. Management of *Rhynchosporium* in Different Barley Varieties and Cropping Systems. London, UK: Home-Grown Cereals Authority, Project Report. 315 pp.
- PEEVER, T.L., ZEIGLER, R.S., DORRANCE, A.E., CORREA-VICTORIA, F.J. and MARTIN, S.S. 2000. Pathogen population genetic and breeding for resistance. ASP net feature. [www.aspnet.org](http://www.aspnet.org)
- ROBBERTSE, B., LENNOX, C.L. VAN JAARVELD, A.B. CROUS, P.W. and VAN DER RIJEST, M. 2000. Pathogenicity of the *Rhynchosporium secalis* population in Western Cape province of South Africa. **Euphytica** 115: 75-82.
- SALAMATI, S. and MAGNUS, H.A. 1997. Leaf blotch on spring barley infected by isolates of *Rhynchosporium secalis* under different temperature and humidity regime. **J. Plant Pathol.** 46: 939-945.
- SALAMATI, S. and TRONSMO, A.M. 1997. Pathogenicity of *Rhynchosporium secalis* isolates from Norway on 30 cultivars of barley. **Plant Pathol.** 46: 416-424.
- SALAMATI, S., ZHAN, J. BURDON, J.J. and McDONALD, B.A. 2000. The genetic structure of field populations of *Rhynchosporium secalis* from three continents suggests moderate gene flow and regular recombination. **Phytopathology** 90: 901-908.

- SHIPTON, W.A., BOYD, W.J.R. and ALI, S.M. 1974. Scald of barley. **Rev. Plant Pathol.** 53: 839-861.
- TEKAUZ, A. 1991. Pathogenic variation in *Rhynchosporium secalis* on barley in Canada. **J. Plant Pathol.** 13: 298-304.
- WILLIAMS, K., DONNELLA, C., SMYLSMOTT, L. and WALWORK, H. 2003. Molecular variation in *Rhynchosporium secalis* isolates obtained from hotspots. **Australas. Plant Pathol.** 37: 257-262.
- XI, K., TURKINGTON, T.K. HELM, J.H. BRIGGS, K.G., TEWARI, J.P.T. and KHARBANDA, P.D. 2003. Distribution of pathotypes of *Rhynchosporium secalis* and cultivar reaction on barley in Alberta. **Plant Dis.** 87: 391-396.
- ZADOKS, J.C., CHANG, T.T. and KONZAK, C.F. 1974. Adecimal code for growth stages of cereals. **Weed Res.** 14: 415-421.
- ZAFFARANO, P.L., McDONALD, B.A. ZALA, M. and LINDE, C.C. 2006. Global hierarchical gene diversity analysis suggests the fertile crescent is not the center of origin of the barley scald pathogen *Rhynchosporium secalis*. **Phytopathology** 96: 941-950.
- ZHAN, J., FITT, B.D.L. PINNSCHMIDT, H.O., OXLEY, S.J.P. and NEWTON, A.C. 2008. Resistance, epidemiology and sustainable management of *Rhynchosporium secalis* population on barley. **Plant Pathol.** 57: 1-14.
- ZHANG, Q., WEBSTER, R.K., CRANDALL, B.A., JACKSON, L.F. and SAGHAI, M.A. 1992. Race composition and pathogenicity associations of *Rhynchosporium secalis* in California. **Phytopathology** 82: 79